

Capturing the diversity of mesoscale trade wind cumuli using complementary approaches from self-supervised deep learning

Dwaipayan Chatterjee¹, Sabrina Schnitt¹, Paula Bigalke¹, Claudia
Acquistapace¹, Susanne Crewell¹

¹Institute for Geophysics and Meteorology, University of Cologne, Cologne, Germany

Key Points:

- Mesoscale cloud organization can be taxonomized by a two-step deep learning approach in the feature space continuum
- Comparison with human-annotated labels reveals the need to include uncertainty estimates of the human-derived classification
- Our approach can describe the temporal transition from human-labeled sugar to flower regimes

Corresponding author: Dwaipayan Chatterjee, dchatter@uni-koeln.de

Abstract

At the mesoscale, trade wind clouds organize with a wide variety of spatial arrangements, which influences their effect on Earth's energy budget. Past studies used high-resolution satellite measurements and clustering/labeling techniques to classify trade wind clouds into distinct classes. However, these methods only capture a part of the observed organization variability. This work proposes an integrated framework using a continuous followed by discrete self-supervised deep learning approach based on cloud optical depth from geostationary satellite measurements. The neural network learns the semantics of cloud system structure and distribution, verified through visualizations of different layers. Our analysis compares classes defined by human labels with machine-identified classes, aiming to address the uncertainties and limitations of both approaches. Additionally, we illustrate a case study of sugar-to-flower transitions, a novel aspect not covered by existing methods.

Plain Language Summary

Clouds are a fundamental player affecting our planet's energy balance, making their accurate representation crucial in climate models. One open question is how they organize on a scale of a few 100 km (mesoscale) within the trade wind region. Satellite observations can help to categorize these clouds, but previous methods had limitations in capturing the full range of cloud arrangements and transitions between different cloud forms. We have introduced a novel approach that utilizes machine learning and geostationary satellite data to address this issue. Our machine learning model autonomously learns to recognize various cloud patterns and distributions. We conducted a comparative analysis between the categories generated by the machine and those identified by human experts to understand the strengths and weaknesses of both methods. Additionally, we explore a case study where clouds undergo a transformation, changing from a structure resembling sugar to one resembling flowers. This particular transformation was found difficult to capture with numerical simulation before. Our approach successfully captures the transition in the machine-learned feature space. Overall, the new approach can help to better understand cloud evolution, which is crucial for improving climate models and predicting how cloud behavior may change in a changing climate.

1 Introduction

Shallow convective clouds are small in individual extent but cover large areas of the tropical oceans, appearing as distinct cloud fields. Due to their radiative and precipitating properties, their representation in climate models is crucial for understanding the current large inter-model spread in predicted cloud feedback and climate sensitivity (Bony & Dufresne, 2005; Nuijens & Siebesma, 2019; Vogel et al., 2022). The EUREC⁴A field campaign (Bony et al., 2017; Stevens et al., 2021), which took place in the North Atlantic Trade (NAT) region around Barbados, aimed at investigating the interplay between clouds, convection, and circulation by deploying a large variety of observations between January and February 2020.

While shallow convection was long seen to produce randomly scattered individual clouds, further understanding has been gathered on the importance of cloud field organization for precipitation (Rauber et al., 2007; Radtke et al., 2022) through cold pool formation and rain evaporation (Seifert et al., 2015; Vogel et al., 2021). Among others, open research questions concern a detailed quantification of the role played by shallow mesoscale cloud organizations in controlling cloud amounts and their radiative response in the trades (Bony et al., 2015; Tomassini et al., 2015; Vogel et al., 2020; Vial et al., 2017).

Introducing four shallow convective organizations (*Sugar*, *Gravel*, *Flower*, *Fish*), with common occurrences on meso-beta (20 to 200 km) and meso-alpha (200 to 2,000

km) scales, Stevens et al. (2020) rely on human-labeled visible satellite images in the NAT region. Sugar clouds consist of small, scattered clouds with a limited vertical extent, while gravel clouds exhibit organized lines or arcs resembling cell-like patterns. Fish clouds display a network resembling fishbones with distinct cloud-free spaces, and flowers represent larger, stratiform cloud structures forming dispersed closed cells. These patterns vary in net cloud radiative feedback (Bony et al., 2020); and, when classified by a deep neural network trained on human-labeled scenes (Rasp et al., 2020), display fundamental differences in cloud fraction and environmental conditions (Schulz et al., 2021). The four patterns exhibit a daily cycle (Vial et al., 2021) and transitions, e.g., from sugar to flower, have been studied in Large-Eddy-Simulation (LES) to identify the governing processes (Narenpitak et al., 2021, 2023; Dauhut et al., 2023).

Yet, imposing four distinct classes on the diversity of the observed organization does not cover the intermediate cloud patterns or transient states, as highlighted by the LES studies. Hence, some dynamic processes important for climate feedback may be ignored or neglected. Also, to our knowledge, there is no description of one of such transitions among different cloud regimes purely using observations. Furthermore, most of the recent studies trying to quantify the labeled well-organized systems find that these four cloud systems occur only around 50% over NAT (Janssens et al., 2021; Schulz et al., 2021; Vial et al., 2021) and have some ambiguities in agreement from the labelers' side (Schulz, 2022). Therefore, to handle such complexity, our first objective is to develop simplified, streamlined representations to effectively understand and capture the entire cloud spectrum's organizational relationships.

There are several different possibilities for ordering the variability of mesoscale cloud systems, such as Janssens et al. (2021) who introduced a set of selected metric spaces for arranging the cloud systems using object-based, scale-based, and retrieved physical-based statistical properties. Utilizing the metric scores and a k-means algorithm, they observe that human-defined classes have better separation starting at seven clusters. Denby (2020) demonstrates that unsupervised neural network models, which involved some human decisions in the learning stage, can be used to distinguish mainly ten different types of cloud organization and their associated radiative properties. In this work, we do not aim to favor any of the presented metrics but rather search for new information purely based on their organizational aspects, minimizing human intervention. Therefore, we aim to identify optimal distinct classes of cloud organizations representing the full spectrum and further compare them with human-identified labels.

Based on GOES-16 E cloud optical depth (COD) images (Sec. 2), Section 3 proposes a two-step self-supervised deep learning approach to study shallow convection in a continuous feature space, characterizing the entire diversity of occurring organizations. Further, an optimized discretization of the continuous space is developed to derive a finite set of classes representative of the continuous spectrum. The representations and their characteristics are investigated in Section 4.1, and we conduct a proof-of-concept study in Section 4.2 to explore the extent of agreement between human-annotated cloud organizations and machine-identified classes. Additionally, we investigate in Section 5 how this approach can be used as a tool to study transitions between different organization patterns.

2 Satellite dataset

We use COD retrieved from GOES-16 E Advanced Baseline Imager (Schmit et al., 2005) by the daytime cloud optical and microphysical properties algorithm (DCOMP) (Walther & Heidinger, 2012) at 2 km horizontal resolution and 10 – 15 minutes temporal resolution. Our domain in NAT (5 – 20° N and 40 – 60° W) is similar to domains used in past studies (Bony et al., 2020; Schulz et al., 2021). The regional climate defines December to May as dry and June to November as wet seasons (Stevens et al., 2016). While

most of the studies focus on dry season shallow convections only, we include some contributions from the wet season by selecting the time period from November to April 2017 – 2021. The purpose of choosing convective occurrences from the wet season is to see how they influence our approach.

COD represents the radiative properties of the cloud in the visible range, and its retrieval from DCOMP tackles the aleatoric uncertainties from the atmosphere and surface robustly. For example, the uncertainty associated with COD retrieval remains below 10% for all ranges in water clouds (see Figure 4 in Walther and Heidinger (2012)). Therefore, we exploit the COD parameter to characterize the cloud system spatio-temporal variability. Note that some fine-scale cloud systems, such as sugar and gravel, also contributing to the variability of mesoscale beta clouds in regional climate systems, may not be fully resolved with the spatial resolution of this product.

Representation learning, also known as feature learning, is a specialized field within machine learning that focuses on extracting meaningful features of a given dataset. To better represent the mesoscale cloud distributions, we use six images per timestamp, including an additional fixed image over the Barbados domain (see S1). Note that the Barbados domain enables comparison with ground-based measurements in future studies. To have an adequate spatial scale of typical occurring cloud fields over NAT (as discussed in Section 1), we use 256 x 256 pixels (roughly 512 square km) as also found in Muller and Held (2012). We exclude crops affected by glint or poor retrieval quality using the respective data flags. Time stamps are limited to 9 am - 3 pm Barbados local time to avoid sun glinting. We utilize land class information to avoid land convection and verify whether 0.98th fraction of random crops belong to the ocean, accepting satellite crops with islands over NAT and excluding those over the northeast South American continent. Finally, to mitigate uncertainties at high COD from DCOMP retrieval, COD values above a threshold of 50, already indicating deep clouds, are clipped to 50. This results in a sample size of 51,000 satellite images.

For further analysis, we make use of hourly ERA-5 (Hersbach et al., 2020) large-scale environmental parameters (horizontal and vertical wind speed, relative humidity) and cloud fraction at a spatial resolution of 0.25°. Hourly cloud amount for four vertical ranges (surface-700 hPa, 700 hPa-500 hPa, 500 hPa-300 hPa, 300 hPa-tropopause) is used from the Clouds and Earth’s Radiant Energy System fourth edition (CERES, Edition - 4A) (Wielicki et al., 1996), characterized by a spatial resolution of 1°.

3 Methods

First, we develop a neural network (N1) that learns to sort the cloud organizations based on the similarity of their visual features, which we call a continuous approach in this work. The purpose is to let the network identify the structural similarities in the cloud systems and map the learned visual features in the 384-dimensional feature space. We use the software package DINO from Facebook Artificial Intelligence Research (FAIR) (Caron et al., 2021) based on PyTorch (Paszke et al., 2019) and the open-source VISSL computer vision library (Goyal et al., 2021) to adapt the network to our requirements. As a backbone neural architecture to process images, we use Vision Transformer (ViT), which has a sequence of self-attention (Vaswani et al., 2023) and feed-forward layers (Bebis & Georgiopoulos, 1994) paralleled with skip connections. This setup helps to identify long-range spatial dependencies by learning relevant information in the image (Khan et al., 2022). To focus on the structural similarities of the cloud system, every epoch, we opt for two random global crops with a 0.75 fraction (192 x 192 pixels) of the parent satellite image. As the largely overlapping global-crop pair has very similar cloud structures, the network learns their essential features and puts the pair closer to each other in the high-dimensional feature space. More details are given in S2.

After obtaining the continuously sorted representation of cloud systems (see Fig. 1.a), we intend to find optimal boundary conditions within the sorted order and, based on it, train a second neural network (N2) to discretize it. As a first step, we reduce the 384-dimension features of the satellite images obtained from N1 to two dimensions using the well-established t-distributed Stochastic Neighbor Embedding (tSNE) algorithm (van der Maaten & Hinton, 2008). tSNE tries to preserve the relative local position between features and the overall global structure of the feature distributions while mapping on a reduced two-dimensional space. On this 2-dimensional representation space, we apply a set of three statistical approaches, namely metric scores of distortion, silhouette (Rousseeuw, 1987), and Calinski-Harabasz (Caliński & Harabasz, 1974) to identify the possible number of optimal classes into which the given features could be clustered. Schubert (2023) suggests taking a collective inference from these three methods to best fit the spherical k-means clustering algorithm used during the training of N2. Supplement 3 illustrates how the three metrics point to an optimal clustering into seven classes.

N2 from Chatterjee et al. (2023) learns to put each satellite image in one of the seven classes as it progressively improves the feature space's clustering, minimizing the cross entropy between two global random crops (192 x 192) from the parent satellite image. Here, the main difference from N1 is that additional augmented image versions (random flipping and noise addition by random Gaussian blur) of global random crops (see Fig. S2.2.b) are included. Augmentations try to provide auxiliary support to the network's generalizability and better capture the differences in diversity of the shallow cloud systems (Nie et al., 2021; Paletta et al., 2023). After obtaining the label of each satellite image, we transfer the assigned class to the continuous representation space, which proves helpful because N1 has learned the sorting arrangement of keeping similar cloud systems closer. Therefore, it helps to visualize how each cluster with distinct characteristics can form a separate local region. The N2 feature space is i) more sparse than N1 (see S2 for explanation) and ii) arranged by closeness to the centroids, which, unlike N1, may not be ideal for representing smooth transitions of cloud systems.

4 Results

4.1 Continuous and discrete representations

To investigate how the satellite images arrange themselves in the feature space of N1, we first study the high-dimensional feature space and assess the arrangement of diverse cloud systems inside it. We reduce the feature dimensions to a 2D space to visualize the continuum using the tSNE algorithm (described in Section 3). Different cloud organizations can be identified in different areas of the 2D space (Fig. 1.a). Going anticlockwise from the top, arch-shaped cloud systems lie in the top-left, followed by flower-type distributions on the left side of the 2D feature space. Close to the flowers in the bottom-left are the flowers spreading out into stratocumulus. Note that while modeling studies suffer from capturing the transition of stratocumulus to cumulus (Sarkar et al., 2020), these cloud regimes are adjacent to one another in the 2D representation.

The bottom part of the feature space contains long bony skeletons, i.e., fish-type cloud systems, and the bottom-right corner shows an extended part of fish-type cloud organizations delineated by unusually large cloud-free regions. The top-right region of the 2D space is a collection of deep convective cells. These primarily occur in the month of November. Arc-shaped cloud systems appear on the left and top-left of the 2-D feature space. Vogel et al. (2021) suggest that the horizontal structure of mesoscale arcs is intrinsically linked to gravel, flowers, and fish. In sequence, Figure 1a shows a continuous link in the spatial arrangement of cloud systems rather than the distinct classes. Additionally, in S4, we investigate how N1 is sensitive to different visual features of cloud organizations and find that the network pays attention to specific patterns in cloud or-




Figure 1. a) Visualization of four hundred randomly selected 256 x 256 satellite images arranged in the dimensionally reduced 2D continuous feature space where the closeness of one satellite image to another is learned by N1. b) Optimized classification learned by N2 provides labels overlaid on the continuous feature space to show the clustering performance. Each class shows low, mid-low, mid-high, and high cloud amounts (%) obtained from the CERES hourly data set. c) Centroid COD images belonging to seven clusters as identified by the discrete neural network (N2). The table shows per class mean of cloud fraction (CF, %) from GOES retrieval and integrated water vapor (IWV, kgm^{-2}) from ERA-5.

ganization, such as deep convective semantics, adjacent thin convection around deep convection, and clear sky features.

Using N2, each of the images can be attributed to one of the seven classes (refer to Section 3), revealing distinct spaces within the 2D continuous representation space (Fig. 1.b). To help investigate how well the seven classes separate, they are evaluated using cloud amount at four different height levels from CERES data. This analysis, on the one hand, reflects how each class differs from the others, and on the other hand, it reasons for the underlying closeness of each class with neighbor classes in the feature space. The difference between the seven clusters is especially evident when looking at their centroid images (Fig. 1.c).

Deep convective class three has by far the highest cloud fraction of 76% and a third more water vapor amount (47.0 kgm^{-2}) than all other classes (mean = 32.5 kgm^{-2}). Neighboring class six (in feature space) includes less frequent higher-level clouds and has a reduced CF of 59% compared to class three. All other classes are dominated by low-level clouds with lower than 50% CF. Classes one and four (neighbor to class six) still have some mid to high-level cloud amount (below 10%). Class one can be interpreted as representing arch-shaped cloud systems, and four resembles the fish class with a more open sky (also shown by reduction in CF). Classes two, five, and seven, being close in the 2D feature space, have similar cloud vertical distributions and IWV ranging from 30 to 32 kgm^{-2} ; however, their organization is very different, as illustrated by the centroids (Fig. 1.c) and mean CFs (43%, 27%, and 33%, respectively). Class two primarily comprises shallow cloud cover, corresponding to cloud systems resembling fish-type formations. Class five has the lowest cloud fraction and is an intermediary class type between classes two and seven. Finally, class seven has a presence of low cloud amounts and negligible mid to higher cloud amounts, which visually resembles flower-type cloud distributions. Therefore, discretizing the continuous feature space helps us visually find three main classes (one, two, and seven) frequently resembling features identified by humans, i.e., sugar, fish, and flower, respectively. However, it also shows the remaining diversity and their characteristics in a cohesive approach.

4.2 Machine versus human labels

While we checked for visual correspondence and class-wise characteristics in Section 4.1, we now aim to quantify how human labels compare to the machine’s seven clusters. We use the seven previously identified cluster boundaries and cloud system positions in the continuous feature space (N1 + N2 together defined as “framework” from now on) and the dataset by Schulz (2022), providing human labels with an agreement score ranging between 0 and 100%.

For each timestamp where at least one of the four patterns was identified within our domain, we select a 256×256 -pixel satellite image centered over the area of highest human agreement. In this way, we ensure the best possible intercomparison. Applying the pre-processing (as in Section 2) leaves us with 52 samples of human-labeled satellite images (fish: 19.3%, gravel: 26.9%, flower: 28.8%, sugar: 25.0%). Note that the best and worst cloud organization agreements with this procedure are 91% and 7%, respectively. Finally, we get the feature vectors of the images corresponding to the human samples from N1 and the machine-identified labels from N2.

The framework classifies 40% flower-labeled cloud systems in class seven (see the hit rate along each class in Fig. 2.a) while sugar-labeled cloud systems are 31% classified in class one and 20% in class four. For class four, note sugar’s low agreement in Fig. 2.b. Gravel has a total of 44% representation in classes one and five, whereas fish annotated labels are allocated 30% in class two and 20% each in classes four and five. Further, looking at example images visually (Fig. 2.a), in contrast to images with high human agreement, it is evident that those with lower agreement significantly deviate from




Figure 2. a) For better visualization and reference purposes of human labels, each column shows 256 x 256 COD images belonging to a certain class marked with the highest and lowest human agreement displayed along the two rows. Below, the images along each column show the proportional machine-predicted class for human labels. b) Continuous feature space colored with different classes (1-7) in the background, along with Human labels (fish, sugar, flower, gravel) in the foreground. The level of human agreement on the identified patterns is indicated by symbol size. c) Relative occurrence of 30 nearest neighbors to human-labeled fish, gravel, flower, and sugar along the seven machine-labeled classes.

the recognized definitions (as given in Stevens et al. (2020)) of sugar, gravel, flower, and fish cloud structures.

Within the 2D feature space (Fig. 2.b), flowers detected with high probability mostly occur in areas of class seven, which was already well reflected in the centroids. Following a similar agreement is sugar (street-type cloud systems), which can be found in areas of class one. However, 38% of sugar samples, with a low agreement, lies in classes four and five, which are extended fish and flower type classes (Section 4.1). Thus, even though these samples reside in those regions of the feature space, their confidence is less than 25%. Rightly, no human-labeled samples are found in class three, which predominantly comprise deep convective cells. For the gravel pattern, 21% samples belong to class six (Fig. 2.b)) and exhibit minimal human confidence; in contrast, the rest from the gravel class are positioned between classes one and seven, suggesting that gravel cloud cell sizes fall between sugar and flower. Finally, the fish class exhibits relatively higher confidence in human labels, aligning well with the feature space characteristics, and lies in class two (fish) and four (extended fish).

To compensate for the limited number of human label samples, we analyze the 30 nearest satellite images to each human label as identified by N1 (Fig. 2.c). This analysis aims to show the generalization capacity of our approach and further enhance our understanding of the connection between organizations. The majority of neighbors in human-identified fish-type cloud systems (more than 50%) belong to machine-identified classes two and four, representing fish and extended fish-type cloud structures with large cloud-free regions. The gravel regime includes members of all classes, with notable contributions from classes one, five, and seven, which exhibit cloud cell characteristics similar to gravel systems. One of the reasons for the wider spread of neighbors might be due to the lower human agreement of the images labeled as gravel (75% of gravel-labeled samples had agreement less than 0.25). In contrast, the flower regime mainly belongs to class seven (46 %), further aligning with the high confidence of human labels. Regarding sugar-type cloud systems, 37 % of the neighbors fall into class one, while those with low human agreement are scattered across the remaining classes. Therefore, we find that machine-labeled classes encompass the human-labeled ones, especially for sugar, flower, and fish, but not so clearly for gravel.

Comparing human labels with their nearest neighbors shows that the framework provides more objective freedom and improves our confidence about the feature vectors allocated to images corresponding to human samples. It also shows the uncertainty associated with less agreed-upon human labels. Further, in S5, using ERA-5 large-scale environmental variables and cloud physical properties, we demonstrate that both the neighbors and the human crops share a similar, homogeneous distribution of physical properties.

5 Transitions

To showcase an application that highlights the strengths and weaknesses of the presented framework, we explore the "sugar" to "flower" (S2F) cloud system transition on February 2, 2020. Using LES, Narenpitak et al. (2021) showed a strengthening of large-scale upward wind motion and an increase in total water path and optical depth as the transformation develops towards the flower. Here, we look at how the transition in COD is represented in the feature space. For example, where do the representations of transitions lie in the feature space? How smooth is the transition in the feature space?

Covering the spatio-temporal developments, 47 COD images were collected (after applying quality filter checks (see Section 2)), centered at 12.5° N, 50° W. They cover the time from 10:50 to 19:20 UTC, with a gap between 17:00 to 18:00 UTC likely caused by local sun glint. We ingest the available samples into the trained framework, collect




Figure 3. a) Five COD images covering the transition period between sugar and flower on the second of February 2020. Their position in the 2D feature space is indicated in the center of the bottom row. b) Individual and standard deviation profiles of 1) vertical, 2) horizontal wind speed describing the atmospheric dynamics, and 3) cloud cover showing changes in mesoscale structure of the transition samples. c) Illustration of temporal transition development inside the feature space: cosine distance of the first daytime image feature obtained at 10:50 UTC compared with the cloud system evolution features for the rest of the day (blue). The last obtained image at 19:20 UTC towards the first image (orange) and θ_m represents the increasing cosine distance.

315 their features (from N1) and machine labels (from N2), and further dimensionally reduce
 316 the features for 2D visualization.

Sugar systems comprise small and shallow clouds with a large spread of individual cloud cells in a domain, as evident in the beginning (10:50, Fig. 3.a). In contrast, flower systems appear in multiple deeper aggregates surrounded by large dry areas and are detected first in the southeast cover at 16:50 before becoming dominated at 19:20 over the full domain. In general, the transition features lie at the border of well-defined clusters one (‘sugar’) and cluster seven (‘flower’) (Fig. 3.a), and the framework is able to capture their intermediary nature as they are neither perfect sugar nor flower type. We use wind speed (vertical and horizontal) to represent changes in atmospheric dynamics and changes in cloud cover to account for the changes in mesoscale structure from the ERA-5 product. A gradual increase in vertical velocity is observed as the system transitions from sugar to flowers, and consequently, the surface wind speed gradually reduces its strength (Fig. 3.b). In addition, as expected, cloud fraction profiles show a gradual decrease as the transition progresses with time.

Sugar-type mesoscale organizations typically occur during the daytime with shallow boundary layers, while flowers occur at night with deeper boundary layers (Vial et al., 2021). We use cosine distance between the features to show the gradual development of the S2F transition inside the feature space (Fig. 3.c), which quantifies the variation in visual features of the 47 COD images. The transformation appears smooth initially, with relatively more significant changes occurring later (post-18:00 UTC) as the system approaches the flower state. We associate the relatively high changes in cosine distance compared to initial sugar stages to convective developments which are faster once the system starts approaching a well-defined flower state. This example illustrates that the framework can capture the intrinsic characteristics of S2F transitions and can be further exploited as a tool to study cloud system transformations and associated processes with large satellite datasets.

6 Conclusion

In this work, we develop and make use of a two-step self-supervised learning approach to study shallow convective organization properties and their transitions. By analyzing organization in a continuous approach without imposing predefined classes, we include all occurring patterns and transitional states in our analysis. Moreover, the approach shows that mesoscale cloud organizations in NAT can be classified into seven optimal classes for the time period considered. Exploiting the cloud amount at different vertical levels from CERES measurements, we show how the classes are interlinked with each other within the continuous space, and thus, the feature space captures the variability of tropical clouds in more detail.

We compare human-labeled cloud systems (Schulz, 2022) to machine-identified cluster regions. Cloud systems with higher agreement among humans lie in the “correct” region of the feature space, while the ones with less consensus are in the “wrong” regions of the feature space. Two of the seven optimal classes are strongly related to flower and sugar, respectively. Representing the sugar-to-flower transition case study (Narenpitak et al., 2021) for February 2, 2020, in the feature space illustrates the capability to identify and represent the observed transformations smoothly in their clearly interpretable regions. We evaluate the transition’s large-scale environmental parameters and observe a gradual increase in vertical wind speed and a gradual decrease in cloud amount. Finally, we demonstrate the framework’s capability to capture the underlying mesoscale visual transformations, such as the transition approaching mature flower convective stages through quick changes in consecutive cosine distances.

One of the limitations of this study is that we use only the daytime cloud retrievals, and hence, the nocturnal nature of the organizations cannot be captured. Future studies will use infrared satellite measurements for 24-hour coverage. We aim to fine-tune our framework with the ground-based observations of the EUREC⁴A campaign and fur-

ther extend our analysis to a climate scale. Currently, Destination Earth (Hoffmann et al., 2023) focuses on simulating high-resolution global digital twins at a 1 km grid scale. The developed workflow could be a testing ground for investigating the newly adjusted subgrid parameterization effects on mesoscale cloud systems or atmospheric processes at different scales.

7 Open Research

CERES, Edition-4A, DOI:10.5067/TERRA+AQUA/CERES/SYN1DEG-1HOUR_L3.004A) is made available by the NASA CERES group. ERA-5 reanalyses were downloaded from the Copernicus climate change services DOI:10.24381/cds.143582cf.

The code to produce this work and pre-trained weights of N1 and N2 can be accessed at <https://doi.org/10.5281/zenodo.8352614>

Acknowledgments

Dwaipayan Chatterjee’s research was supported by the Federal Ministry for Environment, Nature Conservation, Nuclear Safety, and Consumer Protection. Claudia Acquistapace’s (CA) research was funded by Deutsche Forschungsgemeinschaft (DFG). CA also acknowledges funding from Federal Ministry for Digital and Transport (BMDV). The authors would like to thank Andi Walther for providing the GOES-16 E satellite data. This work was performed as part of the Helmholtz School for Data Science in Life, Earth, and Energy (HDS-LEE).

References

- Bebis, G., & Georgiopoulos, M. (1994). Feed-forward neural networks. *IEEE Potentials*, 13(4), 27-31. doi: 10.1109/45.329294
- Bony, S., & Dufresne, J.-L. (2005). Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models. *Geophysical Research Letters*, 32(20). doi: 10.1029/2005GL023851
- Bony, S., Schulz, H., Vial, J., & Stevens, B. (2020). Sugar, Gravel, Fish, and Flowers: Dependence of Mesoscale Patterns of Trade-Wind Clouds on Environmental Conditions. *Geophysical Research Letters*, 47(7), e2019GL085988. doi: 10.1029/2019GL085988
- Bony, S., Stevens, B., Ament, F., Bigorre, S., Chazette, P., Crewell, S., ... Wirth, M. (2017, November). EUREC4A: A Field Campaign to Elucidate the Couplings Between Clouds, Convection and Circulation. *Surveys in Geophysics*, 38(6), 1529–1568. doi: 10.1007/s10712-017-9428-0
- Bony, S., Stevens, B., Frierson, D. M. W., Jakob, C., Kageyama, M., Pincus, R., ... Webb, M. J. (2015). Clouds, circulation and climate sensitivity. *Nature Geoscience*, 8, 261–268. doi: 10.1038/ngeo2398
- Caliński, T., & Harabasz, J. (1974). A dendrite method for cluster analysis. *Communications in Statistics*, 3(1), 1-27. Retrieved from <https://www.tandfonline.com/doi/abs/10.1080/03610927408827101> doi: 10.1080/03610927408827101
- Caron, M., Touvron, H., Misra, I., Jégou, H., Mairal, J., Bojanowski, P., & Joulin, A. (2021). *Emerging properties in self-supervised vision transformers*.
- Chatterjee, D., Acquistapace, C., Deneke, H., & Crewell, S. (2023). Understanding cloud systems structure and organization using a machine’s self-learning approach. *Artificial Intelligence for the Earth Systems*. Retrieved from <https://journals.ametsoc.org/view/journals/aies/aop/AIES-D-22-0096.1/AIES-D-22-0096.1.xml> doi: <https://doi.org/10.1175/AIES-D-22-0096.1>
- Dauhut, T., Couvreur, F., Bouniol, D., Beucher, F., Volkmer, L., Pörtge, V., ...

- Wirth, M. (2023). Flower trade-wind clouds are shallow mesoscale convective systems. *Quarterly Journal of the Royal Meteorological Society*, 149(750), 325–347. doi: 10.1002/qj.4409
- Denby, L. (2020). Discovering the Importance of Mesoscale Cloud Organization Through Unsupervised Classification. *Geophysical Research Letters*, 47(1), e2019GL085190. doi: 10.1029/2019GL085190
- Goyal, P., Duval, Q., Reizenstein, J., Leavitt, M., Xu, M., Lefaudeux, B., ... Misra, I. (2021). *Vissl*. <https://github.com/facebookresearch/vissl>.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... Thépaut, J.-N. (2020). The era5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. doi: <https://doi.org/10.1002/qj.3803>
- Hoffmann, J., Bauer, P., Sandu, I., Wedi, N., Geenen, T., & Thiemert, D. (2023). Destination earth – a digital twin in support of climate services. *Climate Services*, 30, 100394. Retrieved from <https://www.sciencedirect.com/science/article/pii/S2405880723000559> doi: <https://doi.org/10.1016/j.cliser.2023.100394>
- Janssens, M., Vilà-Guerau de Arellano, J., Scheffer, M., Antonissen, C., Siebesma, A. P., & Glassmeier, F. (2021). Cloud Patterns in the Trades Have Four Interpretable Dimensions. *Geophysical Research Letters*, 48(5), e2020GL091001. doi: 10.1029/2020GL091001
- Khan, S., Naseer, M., Hayat, M., Zamir, S. W., Khan, F. S., & Shah, M. (2022, jan). Transformers in vision: A survey. *ACM Computing Surveys*, 54(10s), 1–41. doi: 10.1145/3505244
- Muller, C. J., & Held, I. M. (2012, August). Detailed Investigation of the Self-Aggregation of Convection in Cloud-Resolving Simulations. *Journal of the Atmospheric Sciences*, 69(8), 2551–2565. (Publisher: American Meteorological Society Section: Journal of the Atmospheric Sciences) doi: 10.1175/JAS-D-11-0257.1
- Narenpitak, P., Kazil, J., Yamaguchi, T., Quinn, P., & Feingold, G. (2021). From Sugar to Flowers: A Transition of Shallow Cumulus Organization During ATOMIC. *Journal of Advances in Modeling Earth Systems*, 13(10), e2021MS002619. doi: 10.1029/2021MS002619
- Narenpitak, P., Kazil, J., Yamaguchi, T., Quinn, P. K., & Feingold, G. (2023). The Sugar-To-Flower Shallow Cumulus Transition Under the Influences of Diel Cycle and Free-Tropospheric Mineral Dust. *Journal of Advances in Modeling Earth Systems*, 15(1), e2022MS003228. doi: 10.1029/2022MS003228
- Nie, Y., Zamzam, A. S., & Brandt, A. (2021). Resampling and data augmentation for short-term pv output prediction based on an imbalanced sky images dataset using convolutional neural networks. *Solar Energy*, 224, 341–354. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0038092X21004795> doi: <https://doi.org/10.1016/j.solener.2021.05.095>
- Nuijens, L., & Siebesma, A. P. (2019, June). Boundary Layer Clouds and Convection over Subtropical Oceans in our Current and in a Warmer Climate. *Current Climate Change Reports*, 5(2), 80–94. doi: 10.1007/s40641-019-00126-x
- Paletta, Q., Terrén-Serrano, G., Nie, Y., Li, B., Bieker, J., Zhang, W., ... Feng, C. (2023). Advances in solar forecasting: Computer vision with deep learning. *Advances in Applied Energy*, 11, 100150. Retrieved from <https://www.sciencedirect.com/science/article/pii/S266679242300029X> doi: <https://doi.org/10.1016/j.adapen.2023.100150>
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., ... Chintala, S. (2019). *Pytorch: An imperative style, high-performance deep learning library*.
- Radtke, J., Naumann, A. K., Hagen, M., & Ament, F. (2022). The relationship between precipitation and its spatial pattern in the trades observed during

- EUREC4A. *Quarterly Journal of the Royal Meteorological Society*, 148(745), 1913–1928. doi: 10.1002/qj.4284
- Rasp, S., Schulz, H., Bony, S., & Stevens, B. (2020, November). Combining Crowdsourcing and Deep Learning to Explore the Mesoscale Organization of Shallow Convection. *Bulletin of the American Meteorological Society*, 101(11), E1980–E1995. (Publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society) doi: 10.1175/BAMS-D-19-0324.1
- Rauber, R. M., Stevens, B., Ochs, H. T., Knight, C., Albrecht, B. A., Blyth, A. M., ... Zuidema, P. (2007, December). Rain in Shallow Cumulus Over the Ocean: The RICO Campaign. *Bulletin of the American Meteorological Society*, 88(12), 1912–1928. doi: 10.1175/BAMS-88-12-1912
- Rousseeuw, P. J. (1987). Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *Journal of Computational and Applied Mathematics*, 20, 53–65. Retrieved from <https://www.sciencedirect.com/science/article/pii/0377042787901257> doi: [https://doi.org/10.1016/0377-0427\(87\)90125-7](https://doi.org/10.1016/0377-0427(87)90125-7)
- Sarkar, M., Zuidema, P., Albrecht, B., Ghate, V., Jensen, J., Mohrmann, J., & Wood, R. (2020). Observations pertaining to precipitation within the northeast pacific stratocumulus-to-cumulus transition. *Monthly Weather Review*, 148(3), 1251 – 1273. doi: 10.1175/MWR-D-19-0235.1
- Schmit, T. J., Gunshor, M. M., Menzel, W. P., Gurka, J. J., Li, J., & Bachmeier, A. S. (2005). Introducing the next-generation advanced baseline imager on goes-r. *Bulletin of the American Meteorological Society*, 86(8), 1079 – 1096. Retrieved from <https://journals.ametsoc.org/view/journals/bams/86/8/bams-86-8-1079.xml> doi: <https://doi.org/10.1175/BAMS-86-8-1079>
- Schubert, E. (2023, jun). Stop using the elbow criterion for k-means and how to choose the number of clusters instead. *ACM SIGKDD Explorations Newsletter*, 25(1), 36–42. doi: 10.1145/3606274.3606278
- Schulz, H. (2022). C³ontext: a common consensus on convective organization during the eurec⁴a experiment. *Earth System Science Data*, 14(3), 1233–1256. doi: 10.5194/essd-14-1233-2022
- Schulz, H., Eastman, R., & Stevens, B. (2021). Characterization and Evolution of Organized Shallow Convection in the Downstream North Atlantic Trades. *Journal of Geophysical Research: Atmospheres*, 126(17), e2021JD034575. doi: 10.1029/2021JD034575
- Seifert, A., Heus, T., Pincus, R., & Stevens, B. (2015). Large-eddy simulation of the transient and near-equilibrium behavior of precipitating shallow convection. *Journal of Advances in Modeling Earth Systems*, 7(4), 1918–1937. doi: <https://doi.org/10.1002/2015MS000489>
- Stevens, B., Bony, S., Brogniez, H., Hentgen, L., Hohenegger, C., Kiemle, C., ... Zuidema, P. (2020). Sugar, gravel, fish and flowers: Mesoscale cloud patterns in the trade winds. *Quarterly Journal of the Royal Meteorological Society*, 146(726), 141–152. doi: 10.1002/qj.3662
- Stevens, B., Bony, S., Farrell, D., Ament, F., Blyth, A., Fairall, C., ... Zöger, M. (2021, August). EUREC⁴A. *Earth System Science Data*, 13(8), 4067–4119. (Publisher: Copernicus GmbH) doi: 10.5194/essd-13-4067-2021
- Stevens, B., Farrell, D., Hirsch, L., Jansen, F., Nuijens, L., Serikov, I., ... Prospero, J. M. (2016). The barbados cloud observatory: Anchoring investigations of clouds and circulation on the edge of the itcz. *Bulletin of the American Meteorological Society*, 97(5), 787 – 801. doi: <https://doi.org/10.1175/BAMS-D-14-00247.1>
- Tomassini, L., Voigt, A., & Stevens, B. (2015). On the connection between tropical circulation, convective mixing, and climate sensitivity. *Quarterly Journal of the Royal Meteorological Society*, 141(689), 1404–1416. doi: <https://doi.org/10.1002/qj.2450>

- van der Maaten, L., & Hinton, G. (2008). Visualizing data using t-sne. *Journal of Machine Learning Research*, 9(86), 2579–2605. Retrieved from <http://jmlr.org/papers/v9/vandermaaten08a.html>
- Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A. N., ... Polosukhin, I. (2023). *Attention is all you need*. Retrieved from <https://arxiv.org/abs/1706.03762>
- Vial, J., Bony, S., Stevens, B., & Vogel, R. (2017). Clouds, circulation and climate sensitivity. *Surveys in geophysics*, 38(6), 1331–1353. doi: 10.1007/s10712-017-9418-2
- Vial, J., Vogel, R., & Schulz, H. (2021). On the daily cycle of mesoscale cloud organization in the winter trades. *Quarterly Journal of the Royal Meteorological Society*, 147(738), 2850–2873. doi: 10.1002/qj.4103
- Vogel, R., Albright, A. L., Vial, J., George, G., Stevens, B., & Bony, S. (2022, December). Strong cloud–circulation coupling explains weak trade cumulus feedback. *Nature*, 612(7941), 696–700. (Number: 7941 Publisher: Nature Publishing Group) doi: 10.1038/s41586-022-05364-y
- Vogel, R., Konow, H., Schulz, H., & Zuidema, P. (2021, November). A climatology of trade-wind cumulus cold pools and their link to mesoscale cloud organization. *Atmospheric Chemistry and Physics*, 21(21), 16609–16630. (Publisher: Copernicus GmbH) doi: 10.5194/acp-21-16609-2021
- Vogel, R., Nuijens, L., & Stevens, B. (2020). Influence of deepening and mesoscale organization of shallow convection on stratiform cloudiness in the downstream trades. *Quarterly Journal of the Royal Meteorological Society*, 146(726), 174–185. doi: <https://doi.org/10.1002/qj.3664>
- Walther, A., & Heidinger, A. K. (2012). Implementation of the daytime cloud optical and microphysical properties algorithm (dcomp) in patmos-x. *Journal of Applied Meteorology and Climatology*, 51(7), 1371–1390. doi: 10.1175/JAMC-D-11-0108.1
- Wielicki, B., Barkstrom, B., Harrison, E., Lee, R., Smith, G., & Cooper, J. (1996, 05). Clouds and the earth’s radiant energy system (ceres): An earth observing system experiment. *Bulletin of the American Meteorological Society*, 77, 853–868. doi: 10.1175/1520-0477(1996)077<0853:CATERE>2.0.CO;2

References From the Supporting Information

- Bottou, L. (2012). Stochastic gradient descent tricks. In G. Montavon, G. B. Orr, K.R. Müller (Eds.), *Neural networks: Tricks of the trade: Second edition* (pp. 421–436). Berlin, Heidelberg: Springer Berlin Heidelberg. doi: 10.1007/978-3-642-35289-8_25
- Bridle, J. S. (1989). Probabilistic interpretation of feedforward classification network outputs, with relationships to statistical pattern recognition. In *Nato neurocomputing*. Retrieved from <https://api.semanticscholar.org/CorpusID:59636530>
- Caron, M., Touvron, H., Misra, I., Jégou, H., Mairal, J., Bojanowski, P., Joulin, A. (2021). Emerging properties in self-supervised vision transformers.
- Dosovitskiy, A., Beyer, L., Kolesnikov, A., Weissenborn, D., Zhai, X., Unterthiner, T., Housby, N. (2021). An image is worth 16x16 words: Transformers for image recognition at scale.
- Fukushima, K. (1975). Cognitron: A self-organizing multilayered neural network. *Biological Cybernetics*, 20 (3-4), 121–136. Retrieved 2022-08-30, from <http://link.springer.com/10.1007/BF00342633>doi:10.1007/BF00342633
- Goyal, P., Duval, Q., Reizenstein, J., Leavitt, M., Xu, M., Lefaudeaux, B., Misra, I. (2021). Vissl. <https://github.com/facebookresearch/vissl>

- 575 He, K., Zhang, X., Ren, S., Sun, J. (2015, December). Deep Residual Learning for
576 Image Recognition. Retrieved 2022-08-30, from <http://arxiv.org/abs/1512.03385> (arXiv:1512.03385
577 [cs])
- 578 Hendrycks, D., Gimpel, K. (2023). Gaussian error linear units (gelus). Retrieved
579 from <https://arxiv.org/abs/1606.08415>
- 580 Khan, S., Naseer, M., Hayat, M., Zamir, S. W., Khan, F. S., Shah, M. (2022, jan).
581 Transformers in vision: A survey. *ACM Computing Surveys*, 54 (10s), 1–41. doi: 10.1145/3505244
- 582 Rumelhart, D. E., Hinton, G. E., Williams, R. J. (1986, October). Learning rep-
583 resentations by back-propagating errors. *Nature*, 323 (6088), 533–536. Retrieved from
584 <https://doi.org/10.1038/323533a0> doi: 10.1038/323533a0
- 585 Russakovsky, O., Deng, J., Su, H., Krause, J., Satheesh, S., Ma, S., Fei-Fei, L. (2015,
586 December). ImageNet Large Scale Visual Recognition Challenge. *International Journal*
587 *of Computer Vision*, 115 (3), 211–252. Retrieved 2022-08-31, from [http://link.springer](http://link.springer.com/10.1007/s11263-015-0816-y)
588 [.com/10.1007/s11263-015-0816-y](http://link.springer.com/10.1007/s11263-015-0816-y) doi: 10.1007/s11263-015-0816-y
- 589 Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A. N., . .
590 . Polosukhin, I. (2023). Attention is all you need. Retrieved from [https://arxiv.org/](https://arxiv.org/abs/1706.03762)
591 [abs/1706.03762](https://arxiv.org/abs/1706.03762)